

Macrovipera lebetina obtusa



Common names: West-Asian blunt-nosed viper,^[2] Levant blunt-nosed viper.^[3]

Macrovipera lebetina obtusa is a [venomous viper subspecies^{\[4\]}](#) [endemic](#) to [Asia](#), from central [Turkey](#) to northern [Pakistan](#) ([Kashmir](#)).

Description^[edit]

Scientific Classification

Kingdom: Anamalia
Phylum: Cordata
Class: Reptilia
Order: Squamata
Suborder: Serpentes
Family: Viperidae
Genus Macrovipera
Species: M.lebetina
SubSpecies **M.I.obtusa**

Binomial Name

Macrovipera lebetina obtusa
([Dwigubsky](#), 1832)

Synonyms

- *Vipera obtusa* Dwigubsky, 1832
- *Vipera lebetina obtusa*
— [Terentjev](#) & [Chernov](#), 1940
- *Vipera lebetina obtusa*
— [Minton](#), 1968
- *Daboia (Daboia) lebetina obtusa*
— [Obst](#), 1983
- *Daboia lebetina obtusa*
— [Engelmann](#) et al., 1993
- *Macrovipera l[ebetina]. obtusa*
— [Golay](#) et al., 1993^[1]

Geographic range^[edit]

It is found from central [Turkey](#) through [Syria](#), [Lebanon](#), [Iraq](#), northern [Jordan](#), the [Caucasus](#) region (incl. [Armenia](#)), [Azerbaijan](#), [Dagestan](#), [Iran](#), southern [Afghanistan](#), [Pakistan](#) and the ([Kashmir region](#)). In Pakistan, according to [Khan](#) (1983), *M. lebetina* is restricted to the western highlands; it is [allopatric](#) with [Daboia russelii](#), which occurs in the [Indus River](#) valley.^[5]